

First record of Scymnini (Coleoptera: Coccinellidae), with *Scymnus* (*Pullus*) *rubicundus* Erichson, 1847, in Alagoas, Brazil

DUARTE-DE-MÉLO, Jefferson⁽¹⁾; CAVALCANTI, Joshua Pablo⁽²⁾; CAJÉ, Suianne⁽³⁾; SUÊNIA-BASTOS, Ayane⁽⁴⁾

- ⁽¹⁾  0000-0003-0268-279X. Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba, PR, Brazil. duartedemelojefferson@gmail.com
⁽²⁾  0000-0002-1285-0745. Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba, PR, Brazil. joshuapablo2015@gmail.com.
⁽³⁾  0000-0002-4558-4551. Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba, PR, Brazil. suiannecaje@yahoo.com.
⁽⁴⁾  0000-0002-3443-745X. Setor de Ciências Biológicas, Universidade Federal do Paraná, Curitiba, PR, Brazil. sueniaayane@gmail.com.

The content expressed in this article is the sole responsibility of its authors.

ABSTRACT

Scymnini Mulsant, 1846 (Coleoptera: Coccinellidae) is a cosmopolitan group of inconspicuous beetles, and *Scymnus* Kugelann is the richest genus of this tribe, with more than 700 species in 10 subgenera occurring worldwide, being *Pullus* Mulsant its the most species-rich group. Although all Scymnini genera have been revised for South America, this tribe has not yet been recorded for Alagoas. This article aims to address this gap by providing the first record of *Scymnus* in this state of Brazil, along with an identification key for Scymnini genera in Northeastern Brazil. In January 2025, two specimens of Coccinellidae were collected in Arapiraca, mesoregion of Alagoas, an area characterized by a hot semi-arid climate. The collected beetle is *Scymnus* (*Pullus*) *rubicundus* Erichson, 1847, thus the first record of Scymnini for Alagoas. Records such as these are valuable for our better understanding of the tiny, inconspicuous Coccinellidae, a family of economically important beetles. Knowing this, we also provide an identification key for Scymnini genera from Northeastern Brazil.

RESUMO

Scymnini Mulsant, 1846 (Coleoptera: Coccinellidae) é um grupo cosmopolita de besouros discretos, e *Scymnus* Kugelann é o gênero mais rico desta tribo, com mais de 700 espécies em 10 subgêneros ocorrendo em todo o mundo, sendo *Pullus* Mulsant seu grupo mais diverso. Embora todos os gêneros de Scymnini tenham sido revisados para a América do Sul, esta tribo ainda não foi registrada para Alagoas. Este artigo visa preencher essa lacuna ao apresentar o primeiro registro de *Scymnus* para este estado do Brasil juntamente com uma chave de identificação para os gêneros de Scymnini do Nordeste brasileiro. Em janeiro de 2025, dois espécimes de Coccinellidae foram coletados em Arapiraca, mesorregião de Alagoas, área caracterizada por um clima semiárido quente. O besouro coletado é *Scymnus* (*Pullus*) *rubicundus* Erichson, 1847, portanto o primeiro registro de Scymnini para Alagoas. Registros como este são valiosos para nossa melhor compreensão acerca desses minúsculos e discretos Coccinellidae, uma família de besouros importantes economicamente. Sabendo disso, também fornecemos uma chave de identificação para os gêneros de Scymnini do Nordeste do Brasil.

ARTICLE INFORMATION

Article process:

Submitted: 03/19/2025

Approved: 01/08/2026

Published: 01/10/2026



Keywords:

Identification key,
Lady beetles,
Pullus, *Scymnus*.

Palavras-chave:

Chave de identificação,
Joaninhas,
Pullus, *Scymnus*.

Introduction

Scymnini Mulsant, 1846 (Coleoptera: Coccinellidae) is a cosmopolitan group of inconspicuous lady beetles. Five genera of this tribe are found in South America, and they have all been recently revised for the continent (Gordon, 1996; 2000; González & Gordon, 2002; Gordon, 2007): *Carinoscymnus* Gordon, 2007; *Clitostethus* Weise, 1885; *Nephaspis* Casey, 1899; *Scymnobioides* Casey, 1899; and *Scymnus* Kugelann, 1794.

Scymnus is by far the richest genus of this tribe, with more than 700 species in 10 subgenera. The only subgenus of *Scymnus* naturally occurring in the Neotropics is *Pullus* Mulsant, 1846. *Scymnus* (*Pullus*) is the most speciose group of *Scymnus*, with about 450 species distributed worldwide, 30 species in South America (Gordon, 2000) and 10 species in Brazil (Almeida et al., 2025)—*S. (P.) bennetti* Gordon, 2000; *S. (P.) cerinotum* Gordon, 2000; *S. (P.) demerarensis* Gordon, 2000; *S. (P.) fryi* Gordon, 2000; *S. (P.) gnomus* Gordon, 2000; *S. (P.) kraussi* Gordon, 2000; *S. (P.) loewii* Mulsant, 1850; *S. (P.) murilloi* Gordon, 2000; *S. (P.) rubicundus* Erichson, 1847; and *S. (P.) spanglerorum* Gordon, 2000.

Scymnus is similar to other inconspicuous genera of lady beetles found in Brazil (*Carinoscymnus* Gordon, 2007; *Nephaspis* Casey, 1899; *Scymnus* Kugelann, 1794; *Scymnobioides* Casey, 1899), but can be distinguished from them by the following characters: (1) maxillary apical segment not securiform; (2) prosternal process not projected anteriorly; (3) pronotum without longitudinal carina at margins; (4) antenna without first antennomere expanded; and (5) abdominal postcoxal lines complete. This last aspect may also separate *Scymnus* (*Pullus*) from the exotic *Scymnus* (*Scymnus*), the only other subgenus with record for South America (González et al., 2019).

Despite the revision of all its genera for South America (Gordon, 1996; 2000; González & Gordon, 2002; Gordon, 2007) and recent collection efforts in the State (Oliveira et al., 2022), Scymnini has not yet been recorded for Alagoas. Since the distribution and diversity of species impact political decisions, this paper aims to address this gap by presenting the first record of *Scymnus* for this State of Brazil and providing an identification key for Scymnini genera from Northeastern Brazil.

Material and methods

In January 2025, two adult Coccinellidae were handpicked by the second author JPC in an urban area of the Municipality of Arapiraca (-9.752778, -36.673889; 240 m a.s.l.), mesoregion of the State of Alagoas, Brazil. This area is characterized by a hot semi-arid climate (BSh) according to the Köppen-Geiger climate classification (Kottek et al., 2006; Beck et al., 2023).

The beetles were identified using a key to *Scymnus* (*Pullus*) species of South America (Gordon, 2000). Genitalia were dissected, cleared in 10% KOH, and examined. The material examined was double mounted with the genitalia in microvials and deposited in the Coleção

Entomológica Padre Jesus Santiago Moure (DZUP), Universidade Federal do Paraná, Curitiba, Paraná, Brazil (Curator: Lúcia Massutti de Almeida).

A Leica DMC 2900 digital camera coupled with a Leica M205 C compound stereomicroscope was used to photograph adults and genitalia. Images were edited using Adobe Photoshop CC 2019, version 20.0 (7-day free trial).


The morphological terminology is according to Ślipiński (2007). The classification of Coccinellidae follows Che et al. (2021).

Results and discussion

The specimens consist of two males of *Scymnus (Pullus) rubicundus* Erichson, 1847 (Fig. 1) (DZUP 327497-327498). This species differs from other members of the genus by: (1) lack of maculate pattern; (2) males without tubercle on first ventrite; (3) elytra with at least one row of subsutural punctures; (4) abdominal postcoxal line narrowly separated from posterior margin of ventrite; (5) metasternal punctures about as large as elytral punctures; and (6) dorsal color mostly dark (Gordon, 2000). This is the first record of Scymnini for the State of Alagoas.

Figure 1.

Male of *Scymnus (Pullus) rubicundus* Erichson, 1847 (DZUP 327497), collected in Arapiraca (-9.752778, -36.673889; 240 m a.s.l.), Alagoas, Brazil. (A) Habitus. (B) Abdomen. (C) Tegmen, inner (ventral) view. (D) Tegmen, lateral view. (E) Penis.



Scymnus (Pullus) rubicundus is distributed throughout most of South America, excluding Chile and southern Argentina, and it is the most frequently collected species in this continent (Gordon, 2000), but not even the tribe has been reported for Alagoas, possibly due to insufficient collection efforts or a lack of specialists in the area. However, this species has been recorded from other States in the Northeast Region of Brazil (Table 1). These records are valuable for improving our understanding of beetles that are economically important, including *Scymnus* species.

Table 1.
Distribution by States of the Northeast Region of Brazil and host records of Scymnini Mulsant.

Scymnini species	State(s)	Prey(s)	Plant(s)/Crop(s)	Source(s)
<i>Carinoscymnus bachmanni</i> Gordon, 2007	Maranhão	Not recorded	Not recorded	Gordon, 2007
<i>Nephaspis cocois</i> Gordon, 1972	Pernambuco	<i>Aleurodicus cocois</i> (Curtis, 1846) (Hemiptera: Aleyrodidae)	<i>Anacardium occidentale</i> L. (Anacardiaceae)	Arruda, 1970
<i>Scymnobius pernambucensis</i> Giorgi & González, 2014	Pernambuco	Not recorded	Cotton (Malvaceae)	Giorgi & González, 2014
<i>Scymnus (Pullus) cerinotum</i> Gordon, 2000	Pernambuco (Fernando de Noronha Archipelago)	Not recorded	Not recorded	Rafael et al., 2020
<i>Scymnus (Pullus) demerarensis</i> Gordon, 2000	Bahia	Not recorded	Not recorded	Gordon, 2000
<i>Scymnus (Pullus)</i>	Paraíba	<i>Aphis gossypii</i> Glover, 1877	Cotton (Malvaceae)	Malaquias et al., 2017

<i>rubicundus</i> Erichson, 1847		(Hemiptera: Aphididae)		
	Piauí	<i>Aphis</i> <i>craccivora</i> Koch, 1854 (Hemiptera: Aphididae)	<i>Phaseolus</i> <i>lunatus</i> L. (Fabaceae)	França et al., 2018
	Pernambuco (Fernando de Noronha Archipelago)	Not recorded	Not recorded	Rafael et al., 2020
	Alagoas	Not recorded	Not recorded	This paper

Prey data are scarce, but in general Scymnini are important predators of agricultural pests such as Aleyrodidae and Aphididae (Hemiptera) (Gordon, 1996; 2000). Of the six Scymnini species found in Northeastern Brazil, only *Nephaspis cocois* Gordon, 1972 and *S. (P.) rubicundus* have known host records (Table 1).


Identifying the genera of Scymnini and the tribe itself can be confusing because it includes tiny, inconspicuous beetles that appear very homogeneous to the untrained eye. This tribe can be diagnosed by, among other things: (1) prosternum not expanded anteriorly; (2) apical maxillary palps not securiform; (3) dorsal surface pubescent, barely shiny; and (4) abdomen with six visible ventrites. To aid further research, this work also provides an identification key for the Scymnini genera from Northeastern Brazil.

Identification key to Scymnini genera from Northeastern Brazil

- 1 Pronotum with carinae on lateral margins (Fig. 2A) *Carinoscymnus* Gordon, 2007
- Pronotum without carinae on lateral margins 2
- 2 Antennae with enlarged first antennomere (Fig. 2B) *Nephaspis* Casey, 1899
- Antennae without enlarged first antennomere 3
- 3 Abdominal postcoxal lines complete, recurved (Fig. 1B) *Scymnus* Kugelann, 1794
- Abdominal postcoxal lines incomplete (Fig. 2C) *Scymnobioides* Casey, 1899

Figure 2.

Schematic representation of diagnostic morphological traits of Scymnini genera. (A) Carinoscymnus Gordon, 2007, lateral pronotal carina (arrow). (B) Nephaspis Casey, 1899, first antennomere distinctly enlarged (arrow). (C) Scymnobius Casey, 1899, abdominal postcoxal line (arrow).



Conclusion

This study documents the first occurrence of *Scymnus* (Pullus) *rubicundus* Erichson, 1847 in the State of Alagoas, Brazil, expanding the known distribution of the tribe Scymnini in Northeastern Brazil. The identification of two male specimens, distinguished by unique morphological traits such as the absence of a maculate pattern and a narrowly separated abdominal postcoxal line, highlights the importance of continued entomological surveys in underexplored regions. These findings contribute to a better understanding of the diversity and distribution of economically significant lady beetles in Brazil and provide a foundation for future research and conservation efforts targeting Coccinellidae in semi-arid environments.

Acknowledgements

We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for the fellowships to the first (88887.931875/2024-00), third (88887.931876/2024-00), and fourth (88887.949958/2024-00) authors. We also thank the anonymous reviewers for their valuable comments that improved the initial draft of the manuscript.

Conflict of interest

We have declared that there is no conflict of interest.

REFERENCES

- Almeida, L. M., Churata-Salcedo, J. M., Santos, P. B., Duarte-de-Mélo, J., & Castro, C. F. (2025). *Coccinellidae in Catálogo Taxonômico da Fauna do Brasil*. Available in: <http://fauna.jbrj.gov.br/fauna/faunadobrasil/122061>. Access on 11 March 2025.
- Arruda, E. C. (1970). Nota prévia sobre uma nova espécie do gênero *Nephasis* (Coleoptera, Coccinellidae) predadora do *Aleurodicus cocois* em Pernambuco. *Brazilian Journal of Agriculture - Revista de Agricultura*, 45 (4), 156.
- Beck, H. E., McVicar, T. R., Vergopolan, N., Berg, A., Lutsko, N. J., Dufour, A., Zeng, Z., Jiang, X., van Dijk, A. I. J. M., & Miralles, D. G. (2023). High-resolution (1 km) Köppen-Geiger maps for 1901–2099 based on constrained CMIP6 projections. *Scientific Data*, 10, 724. <https://doi.org/10.1038/s41597-023-02549-6>
- Che, L., Zhang, P., Deng, S., Escalona, H. E., Wang, X., Li, Y., Pang, H., Vandenberg, N., Ślipiński, A., Tomaszewska W. & Liang D. (2021). New insights into the phylogeny and evolution of lady beetles (Coleoptera: Coccinellidae) by extensive sampling of genes and species. *Molecular Phylogenetics and Evolution*, 156, 107045. <https://doi.org/10.1016/j.ympev.2020.107045>
- França, S. M., Melo Júnior, L.C., Gomes Neto, A. V., Ramalho Silva, P. R., Lima, É. F. B. & Melo, J. W. S. (2018). Natural enemies associated with *Phaseolus lunatus* L. (Fabaceae) in Northeast Brazil. *Florida Entomologist*, 101 (4), 688-691. <https://doi.org/10.1653/024.101.0408>
- Giorgi, J. A. & González G. (2014). A new species of *Scymnobi* Casey (Coleoptera, Coccinellidae, Scymnini) from Pernambuco, Brazil. *Revista Brasileira de Entomologia*, 58 (4), 316-318. <https://doi.org/10.1590/s0085-56262014005000008>
- González, G., Lüer, A., & Salinas, R. (2019). Primeros registros de las especies predadoras *Scymnus* (*Scymnus*) *interruptus* (Goeze) y *Scymnus* (*Pullus*) *suturalis* Thunberg (Coleoptera: Coccinellidae) en Chile. *Revista Chilena de Entomología*, 45 (2), 267-272.
- Gordon, R. D. (1996). South American Coccinellidae (Coleoptera). Part V: a taxonomic revision of the genus *Nephaspis* Casey. *Frustula Entomologica*, n.s., 19 (32), 1-50.

- Gordon, R. D. (2000). South American Coccinellidae (Coleoptera). Part VII: A systematic revision of South American *Scymnus* (*Pullus*) Mulsant (Scymninae: Scymnini). *Frustula Entomologica*, n.s., 23 (36): 56-108.
- Gordon, R. D., & González, G. (2002). South American Coccinellidae (Coleoptera). Part IX: A systematic revision of *Scymnobioides* Casey (Scymninae: Scymnini). *Frustula Entomologica*, n.s., 25 (38): 57-85.
- Gordon, R. D. (2007). South American Coccinellidae (Coleoptera). Part XII (lege XIII): *Carinoscymnus*, new genus of South American Scymnini (Scymninae). *Frustula Entomologica* n.s., 30 (43): 161-176.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., & Rubel, F. (2006). World Map of the Köppen-Geiger climate classification updated. *Meteorologische Zeitschrift*, 15, 259-263.
<https://doi.org/10.1127/0941-2948/2006/0130>
- Oliveira, F. P. C., Silva, M. P. B., & Lima, M. S. (2022). Assembleia de coccinelídeos no estado de Alagoas. *Diversitas Journal*, 7 (1), 0190–0202.
<https://doi.org/10.48017/dj.v7i1.1921>
- Malaquias, J. B., Ramalho, F. S., Dias, C. T. S., Brugger, B. P., Lira, A. C. S., Wilcken, C. F., Pachú, J. K. S., & Zanuncio, J. C. (2017). Multivariate approach to quantitative analysis of *Aphis gossypii* Glover (Hemiptera: Aphididae) and their natural enemy populations at different cotton spacings. *Scientific Reports*, 7, 41740.
<https://doi.org/10.1038/srep41740>
- Rafael, J. A., Limeira-de-Oliveira, F., Hutchings, R. W., Miranda, G. F. G., Silva Neto, A. M., Somavilla, A., Camargo, A., Asenjo, A., Pinto. Â. P., Bello, A. M., Dalmorra, C., Mello-Patiu, C. A., Carvalho, C. J. B., Takiya, D. M., Parizotto, D. R., Marques, D. W. A., Cavaleiro, D. O., Mendes, D. M. M., Zeppelini, D., Carneiro, E., Lima, E. F. B., Lima, E. C. A., Godoi, F. S. P., Pessoa, F. A. C., Vaz-de-Mello, F. Z., Sosa-Duque, F. J., Flores, H. F., Fernandes, I. O., Silva-Júnior, J. O., Gomes, L. R. P., Monné, M. L., Castro, M. C. M., Silva, M. P. G., Couri, M. S., Gottschalk, M. S., Soares, M. M. M., Monné, M. A., Rafael, M. S., Casagrande, M. M., Mielke, O. H. H., Grossi, P. C., Pinto, P. J. C., Bartholomay, P. R., Sobral, R., Heleodoro, R. A., Machado, R. J. P., Corrêa, R. C., Hutchings, R. S. G., Ale-Rocha, R., Santos, S. D., Lima, S. P, Mahlmann, T., Silva, V. C. & Fernandes, D. R. R. (2020). Insect (Hexapoda) diversity in the oceanic archipelago of Fernando de Noronha, Brazil: updated taxonomic checklist and new records. *Revista Brasileira de Entomologia*, 64 (3), e20200052.
<https://doi.org/10.1590/1806-9665-RBENT-2020-0052>
- Ślipiński, A. (2007). Australian ladybird beetles (Coleoptera: Coccinellidae). Their biology and classification. Australian Biological Resources Study, Canberra, 286 pp.